

Multiplicity Distributions and Charged-Neutral Fluctuations

Tapan K. Nayak

(for the WA98 Collaboration)

M.M. Aggarwal⁴, A.L.S. Angelis⁷, V. Antonenko¹³, V. Arefiev⁶, V. Astakhov⁶, V. Avdeitchikov⁶, T.C. Awes¹⁶, P.V.K.S. Baba¹⁰, S.K. Badyal¹⁰, S. Bathe¹⁴, B. Batiounia⁶, T. Bernier¹⁵, K.B. Bhalla⁹, V.S. Bhatia⁴, C. Blume¹⁴, D. Bucher¹⁴, H. Büsching¹⁴, L. Carlén¹², S. Chattopadhyay², M.P. Decowski³, H. Delagrange¹⁵, P. Donni⁷, A.K. Dubey¹, M.R. Dutta Majumdar², K. El Chenawi¹², K. Enosawa¹⁸, S. Fokin¹³, V. Frolov⁶, M.S. Ganti², S. Garpman¹², O. Gavrishchuk⁶, F.J.M. Geurts¹⁹, T.K. Ghosh⁸, R. Glasow¹⁴, B. Guskov⁶, H. Å. Gustafsson¹², H. H. Gutbrod⁵, I. Hrivnacova¹⁷, M. Ippolitov¹³, H. Kalechovsky⁷, R. Kamermans¹⁹, K. Karadjev¹³, K. Karpio²⁰, B. W. Kolb⁵, I. Kosarev⁶, I. Koutcheryaev¹³, A. Kugler¹⁷, P. Kulinich³, M. Kurata¹⁸, A. Lebedev¹³, H. Löhrner⁸, L. Luquin¹⁵, D.P. Mahapatra¹, V. Manko¹³, M. Martin⁷, G. Martínez¹⁵, A. Maximov⁶, Y. Miake¹⁸, G.C. Mishra¹, B. Mohanty¹, M.-J. Mora¹⁵, D. Morrison¹¹, T. Mukhanova¹³, D. S. Mukhopadhyay², H. Naef⁷, B. K. Nandi¹, S. K. Nayak¹⁰, T. K. Nayak², A. Nianine¹³, V. Nikitine⁶, S. Nikolaev⁶, P. Nilsson¹², S. Nishimura¹⁸, P. Nomokonov⁶, J. Nystrand¹², A. Oskarsson¹², I. Otterlund¹², T. Peitzmann¹⁴, D. Peressounko¹³, V. Petracek¹⁷, F. Plasil¹⁶, W. Pinganaud¹⁵, M.L. Purschke⁵, J. Rak¹⁷, R. Raniwala⁹, S. Raniwala⁹, N.K. Rao¹⁰, F. Retiere¹⁵, K. Reygers¹⁴, G. Roland³, L. Rosselet⁷, I. Roufanov⁶, C. Roy¹⁵, J.M. Rubio⁷, S.S. Sambyal¹⁰, R. Santo¹⁴, S. Sato¹⁸, H. Schlagheck¹⁴, H.-R. Schmidt⁵, Y. Schutz¹⁵, G. Shabratova⁶, T.H. Shah¹⁰, I. Sibirak¹³, T. Siemiarczuk²⁰, D. Silvermyr¹², B.C. Sinha², N. Slavine⁶, K. Söderström¹², G. Sood⁴, S.P. Sørensen¹¹, P. Stankus¹⁶, G. Stefanek²⁰, P. Steinberg³, E. Stenlund¹², M. Sumner¹⁷, T. Svensson¹², A. Tsvetkov¹³, L. Tykarski²⁰, E.C.v.d. Pijl¹⁹, N.v. Eijndhoven¹⁹, G.J.v. Nieuwenhuizen³, A. Vinogradov¹³, Y.P. Viyogi², A. Vodopianov⁶, S. Vörös⁷, B. Wysłouch³, G.R. Young¹⁶

¹Institute of Physics, 751-005 Bhubaneswar, India ²Variable Energy Cyclotron Centre, Calcutta 700064, India ³MIT Cambridge, MA 02139, USA ⁴University of Panjab, Chandigarh 160014, India ⁵Gesellschaft für Schwerionenforschung (GSI), D-64220 Darmstadt, Germany ⁶Joint Institute for Nuclear Research, RU-141980 Dubna, Russia ⁷University of Geneva, CH-1211 Geneva 4, Switzerland ⁸KVI, University of Groningen, NL-9747 AA Groningen, The Netherlands ⁹University of Rajasthan, Jaipur 302004, Rajasthan, India ¹⁰University of Jammu, Jammu 180001, India ¹¹University of Tennessee, Knoxville, Tennessee 37966, USA ¹²Lund University, SE-221 00 Lund, Sweden ¹³RRC "Kurchatov Institute", RU-123182 Moscow, Russia ¹⁴University of Münster, D-48149 Münster, Germany ¹⁵SUBATECH, Ecole des Mines, Nantes, France ¹⁶Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831-6372, USA ¹⁷Nuclear Physics Institute, CZ-250 68 Rez, Czech Rep. ¹⁸University of Tsukuba, Ibaraki 305, Japan ¹⁹Universiteit Utrecht/NIKHEF, NL-3508 TA Utrecht, The Netherlands ²⁰Institute for Nuclear Studies, 00-681 Warsaw, Poland

Abstract: We discuss the importance of event-by-event fluctuations in relativistic heavy-ion collisions. We present results of multiplicity fluctuations of photons and charged particles produced in 158-A GeV Pb+Pb collisions at CERN-SPS. Multiplicity fluctuations agree with participant model calculations indicating the absence of critical fluctuation. Localized fluctuations in charged particles to photons have been analyzed and compared to various mixed events. The results show localized fluctuations in both charged particles and photons. However, no correlated DCC-like fluctuation is observed.

The primary goal of ultra-relativistic heavy-ion collisions is to investigate the properties of matter at high energy density and high temperature. These studies provide information on the dynamics of multiparticle production mechanism and the phase transition from hadronic matter to Quark-Gluon plasma (QGP). One of the major advantages of

heavy-ion collisions at high energies is that most of the observables can be studied in an event-by-event (E-by-E) basis because of the production of large number of particles in every event. This is particularly important as one expects large fluctuations in several observables associated with the phase transition.

The subject of E-by-E fluctuations has recently gained considerable interest primarily motivated by the near perfect Gaussian distributions of mean transverse momentum and particle ratios [1] obtained by the NA49 experiment at CERN-SPS. As we will see later in this manuscript the multiplicity distributions for narrow centrality bins are also near perfect Gaussians. The variance of the width of these distributions contain information about the reaction mechanism as well as the nuclear geometry. The task here is to distinguish between statistical fluctuations and those which have dynamical origin. Several methods have been put forward which suggest ways to infer about the presence of dynamical fluctuations which have new physics origin. These are done by evaluating the effect of nuclear geometry, influence of hadronic resonances and rescattering, and contributions from Bose-Einstein correlations [2, 3, 4, 5, 6]. Primary interest would be to identify critical fluctuations associated with the QGP phase transition.

Another interesting phenomenon is the formation of disoriented chiral condensates (DCC) [7], which is a consequence of chiral phase transition. It gives rise to large E-by-E fluctuations in the number of charged particles and photons in localized phase space. The distributions of correlation functions using charged particles and photons have been shown to be near perfect Gaussians in the absence of DCC. The presence of DCC tends to make the distributions wider as has been shown in ref. [8].

In order to infer about the presence of non-statistical fluctuations, one has to compare the distributions obtained from experimental data to known models which incorporate all the known phenomena. An alternate or may be complimentary procedure to probe these fluctuations in a model independent manner would be to compare data distributions to those of mixed events generated from the data. Once properly understood, mixed events provide the most unambiguous means to probe any new physics. In this manuscript we present results from Pb+Pb collisions at 158 · AGeV taken at CERN-SPS by the WA98 experimental setup. The results will be compared to a participant model and also to several different types of mixed events.

In the WA98 experiment, the emphasis has been on high precision, simultaneous detection of both photons and hadrons. Photon multiplicities ($N_{\gamma\text{-like}}$) were measured by the preshower photon multiplicity detector (PMD) placed at 21.4 meters from the target and covering an η range of 2.9 – 4.2. Charged particle hits (N_{ch}) were counted using a circular silicon pad detector (SPMD) located 32.8 cm from the target with a coverage of $2.35 < \eta < 3.75$. The centrality of the interaction is determined by the total transverse energy (E_T) measured in the mid rapidity calorimeter. The centralities are expressed as fractions of the measured cross section from the minimum bias E_T distributions.

Figure 1 shows the minimum bias distributions of $N_{\gamma\text{-like}}$ clusters and N_{ch} . Superimposed to these are the distributions corresponding to the centrality cuts of top 1%, 2% and 5% of the minimum bias E_T distributions. These distributions turn out to be near perfect Gaussians with χ^2/ndf close to unity. The solid lines show the Gaussian fits to these curves. It has been observed that the distributions deviate from Gaussians if the centrality bin is made broader than 0 – 5%. In order to study the centrality dependence of fluctuations we have chosen 2% bins in centrality, viz., 0-2%, 2-4%, ..., 58-60%, ... etc. The resulting multiplicity distributions are near perfect Gaussians over large centrality range, from peripheral to central collisions.

If the distribution of a quantity, X , is Gaussian, then the amount of fluctuation may be defined as:

$$\omega_X = \frac{\sigma_X^2}{\langle X \rangle}, \quad (1)$$

where σ_X is the variance of the distribution. and $\langle X \rangle$ denotes the mean value of X . In order to compare the fluctuation for peripheral to central collisions, we have calculated the relative fluctuation defined as ω_X per number of participant nuclei, N_{part} (obtained from VENUS event generator).

Relative multiplicity fluctuations of photons ($\omega_\gamma/N_{\text{part}}$) and charged particles ($\omega_{\text{ch}}/N_{\text{part}}$), as functions of N_{part} are shown in left and right panels of Figure 2, respectively. Since the measured data for photons contain contaminations from charged particles, these have been corrected to obtain distributions for photons only by using the efficiency and purity of the measured photon-like clusters [9]. The resulting data values have been compared with simple calculations based on participant model [6, 10] and also those obtained from VENUS event generator. A reasonable agreement have been seen with the participant model as well as VENUS indicating the absence of any critical fluctuations.

Theoretical predictions suggest that the isospin fluctuations caused by DCC would produce clusters of coherent pions in phase space forming domains localized in phase space. We present results based on E-by-E fluctuation in the relative number of charged particles and photons. Details may be obtained from Ref. [11]. Two different methods are used for DCC search. The first method deals with the correlation of $N_{\gamma\text{-like}}$ and N_{ch} and the second method is based on the discrete wavelet techniques (DWT).




FIGURE 1. Minimum bias distributions of (a) γ -like clusters and (b) charged particle multiplicity produced in Pb induced reactions at 158-A GeV on Pb. Superimposed are the multiplicity distributions for the top 1%, 2%, and 5% most central events. These distributions turn out to be near perfect Gaussians as seen by the solid fitted lines.




FIGURE 2. The relative fluctuations of (a) photons and (b) the charged particles compared to calculations from a participant model and VENUS. The photon data have been corrected from $N_{\gamma\text{-like}}$ to N_{γ} using the efficiency and purity of the measured photon sample. Reasonable agreement has been seen between data and models.

The correlation of $N_{\gamma\text{-like}}$ and N_{ch} has been studied in smaller ϕ -segments by dividing the ϕ -space into 2,4,8 and 16 bins. A common correlation axis (Z) has been obtained by fitting the distribution of the correlations with a second order polynomial. The distribution of the closest distance of the data points to the correlation axis represents the relative fluctuation of $N_{\gamma\text{-like}}$ and N_{ch} for any given ϕ bin. The final distributions (S_Z) for 4 and 8 bins are shown in the left-top panel of Figure 3. Superimposed are the distributions from GEANT simulated events and a particular set of mixed events, called M1.

We have also used the DWT method where the analysis is performed by dividing the azimuthal space of PMD and SPMD into 2^j bins where j is called the scale of the division. The input to the DWT analysis is a spectrum of the function, $N_{\gamma\text{-like}} / (N_{\gamma\text{-like}} + N_{\text{ch}})$, at the highest resolution scale, $j_{\text{max}} (= 5)$. The output consists of a set of wavelet or the

TABLE 1. Four different types of mixed events are constructed by combinations of PMD and SPMD hits. The last column provides the usefulness of each of the mixed events to probe different type of fluctuation.

	PMD	SPMD	Type of fluctuation to probe:
M1	Mix hits	Mix hits	Correlated + Individual
M2	Unaltered hits	Unaltered hits	Correlated
M3- γ	Unaltered hits	Mix hits	N_γ only
M3-ch	Mix hits	Unaltered hits	N_{ch} only

so called father function coefficients (FFC) at each scale, from $j = 1$ to $j = j_{\max} - 1$. The FFC distributions at 4 and 8 bins in ϕ are shown in the left-bottom panel of Figure 3. The S_Z and FFC distributions are Gaussian in nature in the absence of DCC. The presence of events with DCC domains of a particular bin size in ϕ would result in a broader distribution of S_Z and FFC distributions compared to normal events [8, 11].

The presence of DCC-like fluctuations in the experimental data can be inferred by comparing the data distributions to different sets of mixed events. Properly constructed mixed events preserve all the detector effects while removing correlations. Four sets of mixed events are generated by maintaining the global (bin 1) $N_{\gamma\text{like}}-N_{ch}$ correlations as in the real data. Table 1 summarizes the construction of mixed events and the type of fluctuation these can probe. These behavior has been understood by simulating the effect of DCC and analyzing the simulated data in a manner identical to the real data.




FIGURE 3. The S_Z and FFC distributions for 4 and 8 divisions in ϕ are shown in the left panel. The experimental data, M1 and V+G events are shown by solid circles, solid histograms and dashed histograms, respectively. The statistics for data and mixed events are the same, whereas the distribution for the V+G events is normalized to the number of data events. The right panel shows the root mean square (rms) deviations of the S_Z and FFC distributions for various divisions in the azimuthal angle.

The rms deviations of the S_Z and FFC distributions as a function of the number of bins in azimuth are shown for experimental data, four types of mixed events and Geant simulated data in the right panel of Figure 3. The statistical errors are small and the systematic errors are shown in the figure.

First we compare the rms deviations obtained from the data to those of M1 mixed events. The data points are much larger compared to those of M1 for both S_Z and FFC distributions. For 2, 4 and 8 bins the values of the S_Z rms deviations

of data are 2.5σ , 3.0σ and 2.4σ larger than those of the M1 events. Similarly for 4 and 8 bins the values of the FFC rms deviations of data are 3.7σ and 2.8σ larger than those of the M1 events. These indicate the presence of localized non-statistical fluctuations. The observed effect may arise because of (a) E-by-E correlated $N_{\gamma\text{-like}}\text{-}N_{\text{ch}}$ fluctuations, (b) fluctuations in $N_{\gamma\text{-like}}$, and (c) fluctuations in N_{ch} . The source of the fluctuation can be obtained by comparing the data with M2 and M3 types of events.

Next we compare data rms deviations to those of M2. It is observed that in all cases the results from M2 match completely with data. This implies the absence of any E-by-E correlated (DCC-like) fluctuations and allows us only to put an upper limit on DCC production [11].

Presence of individual fluctuations are to be inferred by comparing data with M3 type of events. The M3 type of events are found to be similar to each other within quoted errors and lie between M1 and M2. This indicates presence of localized independent fluctuations in both $N_{\gamma\text{-like}}$ and N_{ch} .

In summary, we have presented multiplicity distributions and multiplicity fluctuations of photons and charged particles produced in 158-A GeV Pb+Pb collisions at the CERN-SPS. Multiplicity fluctuations from data agree reasonably well with participant model calculations over a wide range of centrality, from peripheral to central collisions. This indicates the absence of any critical fluctuations indicative of phase transition at SPS energies. Localized fluctuations in photons and charged particles have been studied in order to search for formation of domains of DCC. The results from correlation and DWT analysis are compared to those of mixed events generated from data. The results provide model independent evidence for non-statistical fluctuations in the data for ϕ interval between 45 and 90 degrees and one unit in pseudorapidity. This is seen to be due to individual fluctuations in both photons and charged particles. No significant E-by-E correlated fluctuations are observed, contrary to expectations for a DCC effect.

REFERENCES

1. H. Appelshauser et al., (NA49 Collaboration), Phys. Lett. **B459** (1999) 679.
2. S. Mrowczynski, Phys. Lett. **B459** 13 (1999); M. Gazdzicki and S. Mrowczynski, Z. Phys. **C54** 127 (1992).
3. M. Stephanov, K. Rajagopal and E. Shuryak, Phys. Rev. Lett. **81** 4816 (1998).
4. M. Asakawa, U. Heinz, and B. Müller, Phys. Rev. Lett. **85** 2072 (2000).
5. S. Jeon and V. Koch, Phys. Rev. Lett. **85**, 2076 (2000).
6. G. Baym and H. Heiselberg, Phys. Lett. **B469** 7 (1999).
7. J.D. Bjorken, Int. J. Mod. Phys. **A7** (1992) 4189; K. Rajagopal and F. Wilczek, Nucl. Phys. **B399** 395 (1993); J. -P. Blaizot and A. Krzywicki, Phys. Rev. **D46** 246 (1992).
8. B.K. Nandi, T.K. Nayak, B. Mohanty, D.P. Mahapatra and Y.P. Vayogi, Phys. Lett. **B461** 142 (1999).
9. M.M. Aggarwal et al., (WA98 Collaboration), Phys. Lett. B **458** 422 (1999).
10. M.M. Aggarwal et al., (WA98 Collaboration), under preparation.
11. M.M. Aggarwal et al., (WA98 Collaboration) Phys. Rev. **C64** 011901R (2001).